



**abomin**  
NATURAL MINERAL FERTILISER



**OLIVE**

Certified Organic by



Certified Organic by



## THE ADVANTAGES AND OBJECTIVES OF USING ABOMIN

- ✓ Increased photosynthesis and fertility
- ✓ Adverse weather resistance
- ✓ Production of oleic acid
- ✓ Resistance to bacterial attack

## DOSAGE AND METHOD OF USE

It is possible to use Abomin in a single type or also to combine the three types of product together (Abomin, Abomin Complex, Abomin Amino).



### BASIC APPLICATION

#### I<sup>a</sup> APPLICATION

(30-60 days after harvest)

300 Kg per Ha

500 Kg per Ha

350 Kg per Ha

#### II<sup>a</sup> APPLICATION

(30 days before flowering)

300 Kg per Ha

500 Kg per Ha

350 Kg per Ha

Apply along the row (70 cm left and right from the row)

### FOLIAR APPLICATION

#### 4-5 APPLICATIONS

(pre and post-flowering)

30 Kg

Apply with water

### IDEAL FOR

Land with > 1.5% organic substance

Also land with organic substance < 1.5%

Accelerates the Abomin process

For foliar application on all cultivations

## COMPOSITION

ELEMENTS	PPM	ELEMENTS	PPM	ELEMENTS	PPM	ELEMENTS	PPM
Potassium (K)	7200	Cobalt (Co)	31	Copper (Cu)	48.74	Samarium (Sm)	8
Calcium (Ca)	56200	Selenium (Se)	0.4	Nickel (Ni)	5.3	Germanium (Ge)	1.1
Magnesium (Mg)	16200	Lithium (Li)	9.6	Molybdenum (Mo)	0.4	Vanadium (Vn)	5.2
Phosphorus (P)	3735	Lanthanum (La)	74.42	Silicon (Si)	3824	Praseodymium (Pr)	21
Iron (Fe)	3952	Cerium (Ce)	181	Strontium (Sr)	249	Silver (Ag)	0.3
Boron (B)	1.29	Barium (Ba)	34	Yttrium (Y)	1.7	Caesium (Cs)	0.85
Manganese (Mn)	1,160.8	Rubidium (Rb)	28	Bismuth (Bi)	0.03	Titanium (Ti)	0.46
Zinc (Zn)	119	Sulphur (S)	200	Base saturation (V)	94.96%	pH	7.8